

MAINSAIL™ WDG

A WATER DISPERSIBLE GRANULAR FUNGICIDE CONTAINING CHLOROTHALONIL FOR SPRAY APPLICATION TO TURF AND ORNAMENTALS.

Mainsail WDG is a contact fungicide that provides broad-spectrum control of 14 turfgrass diseases and algae, over 40 diseases of field grown and container ornamentals, and numerous tree and orchard crop diseases. Effective for strategic use in programs that attempt to minimize disease resistance to fungicides.

Active Ingredient

Chlorothalonil 82.5%

Packaging

- 4 x 5 LB jugs / CS
- 4 x 10 LB jugs / CS

Application

Do not apply within 150 ft. (for air-blast) or 25 ft. (for ground applications) of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the body of water.

Turfgrass – For best results, use as a preventive application when conditions are favorable for disease development. Under severe disease conditions, use the highest rate and shortest interval corresponding with the application schedule. Apply Mainsail WDG with an adequate amount of water to provide complete coverage. Do not mow or water after treatment until spray deposited on turfgrass is thoroughly dry. Mainsail WDG should always be used in conjunction with good turf management practices.

Use of this chemical on home lawns is prohibited.

Sod farm turf treated with chlorothalonil prior to harvest must be mechanically cut, rolled, and harvested.

Ornamentals – For best results, use as a preventive application when conditions are favorable for disease development. Apply Mainsail WDG with an adequate amount of water (until run-off occurs) to provide complete coverage. Repeat applications at 7 to 14 day intervals until conditions are no longer favorable for disease development. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, apply at 7 day intervals. Apply to plants when foliage and flowers are dry, or nearly dry.

Do not use mist blowers or high-pressure spray equipment when making applications in greenhouses.

The user should test for possible phytotoxic responses on ornamental plants, using recommended rates on a small area prior to commercial use. Applications made during bloom may damage flowers and/or fruits.

Fruits and other structures, which may be borne on treated plants, must NOT be eaten.

MAINSAIL™ WDG

Application Rates

Turf – 1 to 5 oz per 1,000 square feet on 7 to 14 day intervals, depending on the disease. See complete label for specific application rates and intervals, and for snow mold and algae control recommendations.

Ornamentals – Apply at a rate of 1.4 pounds per 100 gallons of water on 7 to 14 day intervals unless other directions are given on the label.

Re-entry Intervals (REIs)

Agricultural Use Requirements: 12 hours (Refer to label for special eye irritation provisions.)

Non-Agricultural Use Requirements: Do not allow entrance to treated areas until spray has dried.

Benefits

- Broad spectrum fungicide for the control of numerous turf, ornamental, and tree and orchard crop diseases
- Contains chlorothalonil for very effective and economical control of plant diseases
- Labeled for use on a broad range of turfgrass, ornamental, and tree and orchard plant species
- Effective for strategic use in disease resistance management programs
- High-quality, water dispersible granular formulation for easy and convenient spray application
- Conveniently packaged in 4 x 5 pound jugs and 4 x 10 pound jugs per case

Important

Always read label thoroughly before application. Mainsail WDG, like all dry, sprayable formulations, should be agitated continuously during the application.

Refer to the label for other information and precautions as well as directions for use and disposal.