

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING ►



PROXY®

Growth Regulator

For Commercial Use on Turf including Golf, Parks, and Sports Turf.

Not For Residential Use.

ACTIVE INGREDIENT:

Ethephon [(2-chloroethyl)phosphonic acid]* 21.7%

OTHER INGREDIENTS: 78.3%

*Contains 2 pounds ethephon per gallon.

EPA Reg. No. 432-1230

**KEEP OUT OF REACH OF CHILDREN
DANGER • PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL And TRANSPORTATION Emergencies
ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

Net Contents
2.5 Gallons

4249932

4587404 050426A3

BACKED
by **BAYER™**

FIRST AID

IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or ambulance, and then give artificial respiration preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. No specific antidote is available. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

Victims of severe overexposure, by inhalation, should be kept under medical observation for up to 72 hours for delayed onset of pulmonary edema. In a victim of overexposure by ingestion, careful gastric lavage is required due to the possibility of stomach or esophageal perforation. This material is an acid, but the use of alkaline substances to neutralize it is contraindicated.

PRECAUTIONARY STATEMENTS

DANGER

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive: Causes irreversible eye damage. Wear safety goggles when handling. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Do not inhale vapors, as this product will irritate mucous membranes.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance selection chart.

Applicators and other handlers must wear long-sleeved shirts and long pants, chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes plus socks and protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove clothing immediately if pesticide gets inside. Then wash body thoroughly and change into clean clothing. The contaminated clothing should be washed before reuse.

Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations. Read entire label before using this product.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE

Store in original container and keep tightly closed. Store in a cool dry place.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Offer for reconditioning, if appropriate. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL USE

Proxy[®] Growth Regulator is a plant growth regulator that may be used to suppress the formation of seedheads of various plants including *Poa annua* and white clover and also to suppress the growth of certain cool season grasses. PROXY Growth Regulator is foliarly absorbed and is most effective on actively growing healthy turf. For best results, apply in sufficient volume of water to provide uniform coverage. Use of a spreader/sticker with an application of PROXY Growth Regulator is not necessary. PROXY Growth Regulator is rainfast within 2 hours. Do not allow entry into treated area until PROXY Growth Regulator has dried. For maximum performance, delay mowing until the day after application.

PRECAUTIONS AND RESTRICTIONS

Do not use this product for purposes other than those listed on the label.

Do not treat turfgrass with poor root systems or growing under stress due to poor soil conditions, drought, disease, or insect damage.

Do not use PROXY Growth Regulator in areas where excessive thatch has accumulated.

Do not contaminate water used for irrigation or domestic purposes.

Do not apply PROXY Growth Regulator through any type of irrigation system.

Avoid spray drift to nearby crops, as this product will cause modifications in plant growth. Plant injury may result.

Scalping may occur on creeping bentgrass cultivars after more than 2 applications of Proxy Growth Regulator for *Poa* seedhead suppression.

Proxy Growth Regulator has been used successfully on many bentgrass cultivars. Tolerance testing should be done on new cultivars before extensive use.

Mix only the amount of spray you expect to use each day. Do not allow mixed solution to stand overnight.

Do not buffer spray solutions.

Test tank mixes with other products on a small area before using widely.

IMMEDIATELY RINSE ANY SPILLS WITH PLENTY OF WATER AS PROXY GROWTH REGULATOR IS CORROSIVE.

DO NOT MIX PROXY GROWTH REGULATOR WITH AMMONIUM THIOSULFATE. SUCH TANK MIXTURES MAY RESULT IN FORMATION OF TOXIC FUMES

DO NOT EXCEED A MAXIMUM OF 30 OZ OF PROXY GROWTH REGULATOR PER 1,000 SQ FT PER YEAR.

EQUIPMENT CLEANING

Prolonged exposure to spray deposits will damage acrylic plastics, certain paints, and metals because of the acidic nature of this product.

Rinse thoroughly all exposed acrylic-plastic materials and painted surfaces with a detergent and water within one hour after exposure to spray deposits.

TURF

Seedhead Suppression: Foliar application of Proxy Growth Regulator will provide suppression of *Poa annua* and white clover seedheads. Make the initial application prior to the emergence of new seedheads. A period of 2-3 weeks after application is required for maximum performance. Repeat applications may be made to predominately *Poa annua* or white clover sites as needed, but not less than two weeks after the previous application. Follow other general recommendations and precautions on the label.

APPLICATION	SITES	RATE	SPRAY VOLUME
<u><i>Poa annua</i> and White Clover Seedhead Suppression:</u> Reapplication interval: 2 weeks or greater for all labeled grasses	Golf course turf including Greens, Tees, Fairways, and Roughs Commercial Turfgrasses including Bentgrass, Kentucky Bluegrass, Perennial Ryegrass, Tall & Fine Fescue, and Bermudagrass	5 fl oz/1,000 sq ft	1.0-2.0 gal/1,000 sq ft

Tank Mixture with Primo MAXX: Tank mixtures of Proxy Growth Regulator at 5 oz per 1,000 sq ft and Primo MAXX at 0.125 or 0.25 ounce per 1,000 sq ft can be used to promote seedhead suppression as well as turfgrass quality. Multiple applications of the tank mix combination may be needed. The number of applications must not exceed recommendations for rates or timings for either

product applied once. Follow the recommendation on each product label for the most restrictive application interval for each turfgrass. Application of this tank mixture during frost periods may cause temporary turf discoloration.

Growth Suppression: An application of PROXY Growth Regulator slows the growth of turfgrasses, thus reducing the required frequency of mowing and the volume of clippings collected. For best turfgrass growth regulation, apply Proxy Growth Regulator when daytime air temperatures are 65°F and rising. Proxy Growth Regulator should only be applied once turfgrass mowing heights have been established for the season. Avoid multiple applications of Proxy Growth Regulator in areas where excessive thatch has accumulated.

APPLICATION	SITES	RATE	SPRAY VOLUME
<p><u>Turfgrass Growth Regulation</u> Reapplication intervals:</p> <ul style="list-style-type: none"> • Kentucky Bluegrass-7 weeks • Perennial Ryegrass-7 weeks • Tall/Fine Fescue-4 weeks • Bentgrass-4 weeks 	<p>Golf course turf including Greens, Tees, Fairways, and Roughs</p> <p>Commercial Turfgrasses including Bentgrass, Kentucky Bluegrass, Perennial Ryegrass, Tall & Fine Fescue</p>	<p>5 fl oz per 1,000 sq ft</p>	<p>1.0-2.0 gal per 1,000 sq ft</p>

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd), PROXY® and Backed by Bayer™ are trademarks of Bayer.



Produced for:

Bayer Environmental Science

A Business Group of Bayer CropScience LP

2 T. W. Alexander Drive

Research Triangle Park, NC 27709