

Home & Garden
DIPEL[®]
DUST
CONTAINS DIPEL[®]
BIOLOGICAL INSECTICIDE

ACTIVE INGREDIENTS:

Bacillus thuringiensis Berliner,
 Var. Kurstaki potency of 200
 International Units per milligram*...0.04%

INERT INGREDIENTS:99.96%

TOTAL.....100.00%

*Equivalent to 90 Million International Units per pound.

DIPEL[®] is a Trademark of Abbott Laboratories.

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC
ANIMALS

CAUTION: May cause irritation of eyes, nose, throat, and skin. Avoid inhalation. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, and ponds. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool dry area. Do not store for extended periods of time or where temperatures are above 90°F.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

EPA EST. NO.829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5)

Number following Est. No. corresponds to first digit in Lot No. on bottom of bag.

EPA REG. No. 829-196

DIRECTIONS FOR USE

It is a violation Federal Law to use this product in a manner inconsistent with its labeling.

Apply recommended amounts when air is calm using a hand duster. Apply at first sign of infestation and repeat at weekly intervals as needed to maintain control. To be effective Dipel Dust must be eaten by the insect. Thorough leaf coverage, especially to the underside of the leaves is essential. Some insects stop feeding after application and appear to be unaffected for several days after application. Check foliage for signs of fresh feeding if insects appear to remain active after applying. Apply higher dosage to maturing plants. Do not exceed maximum dosage, nor use less than the lower dosage.

CROP	PEST	OZS. PER 1,500 SQ.FT.
BROCCOLI	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
CABBAGE	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
CAULIFLOWER	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
CELERY	Cabbage Looper	11 - 33
	Cabbage Looper	11 - 33
COLLARDS	Imported Cabbageworm	9 - 27 1/2
	Cabbage Looper	9 - 27 1/2
CUCUMBERS	Cabbage Looper	9 - 27 1/2
	Grape Leaf Folder	9 - 27 1/2
GRAPES	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
KALE	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
LETTUCE	Cabbage Looper	11 - 33
	Cabbage Looper	9 - 27 1/2
MELONS	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
MUSTARD GREENS	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
POTATOES	Cabbage Looper	11 - 33
	Cabbage Looper	11 - 33
TOMATOES	Tomato Hornworm	9 - 27 1/2
	Cabbage Looper	11 - 33
TURNIP GREENS	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2

NOTE: Start treating as soon as possible after hatching and before larvae are protected by leaf folds. Apply to obtain thorough coverage of leaf surfaces.

Dipel Dust may be applied at the recommended rate up to the day of Harvest.

The manufacturer makes no warranties, expressed or implied, concerning this product or the results obtained from its use which extend beyond the description and precautions given herewith.

EPA REG. No. 829-196



Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28793

Boone, NC 28607

revised 1998